

Parametricity and excluded middle

Auke Booij

University of Birmingham

Abstract

In univalent foundations, it is known that the law of excluded middle allows one to define a family of functions $f_X : X \rightarrow X$ that is not the identity function on the booleans. We show that the converse holds as well: given such a function, we derive the law of excluded middle.

Suppose we are given a polymorphic function

$$f_X : X \rightarrow X,$$

where $X : \mathcal{U}$ is its type parameter.

If this were a term in a language such as System F, then parametricity tells us that it must be equal to the identity function id_X for every type X . But parametricity is a metatheoretical framework: it gives properties about the terms of a language, rather than internally stating properties of elements.

Internal to univalent foundations, if we have LEM, then there exists a polymorphic function f such that $f_{\mathbf{2}}$ (where $\mathbf{2}$ is the type of booleans) is not the identity function [2, exercise 6.9]. Since LEM is consistent with univalent foundations, this means that there cannot be an internal proof that a polymorphic function $f_X : X \rightarrow X$ is equal to the identity.

We prove that, in univalent foundations, LEM is precisely what is needed to get a function family not equal to the identity on $\mathbf{2}$: on the one hand, we already know that LEM gives us such a function; on the other hand, we have the following converse.

Theorem 1. *If there is a function $f : \Pi_{X:\mathcal{U}} X \rightarrow X$ with $f_{\mathbf{2}} \neq \text{id}_{\mathbf{2}}$, then LEM holds.*

Alternatively, to confine the amount of univalence needed, we can work in the setting of intensional type theory with function extensionality (but without full univalence), and assume that f is *extensional* in the sense that it is invariant under equivalences on the type X it acts on.

The idea of the proof is that we define a type $\mathbf{3}_P$, which, depending on whether P holds, may or may not be equivalent to $\mathbf{2}$. We then evaluate f at the type $\mathbf{3}_P \simeq \mathbf{3}_P$ (rather than $\mathbf{3}_P$ itself), and prove $P + \neg P$ using that evaluation.

This proof has been formalized [1] in Agda using the HoTT library.

Proof. Without loss of generality, we may assume that $f_{\mathbf{2}}(0_{\mathbf{2}}) \neq 0_{\mathbf{2}}$.

To prove LEM, let P be an arbitrary proposition. We need to prove $P + \neg P$.

We will consider a type with three points, where we identify two points depending on whether P holds. Formally, this is the quotient of a three-element type, where the relation between two of those points is the proposition P . This quotient can be constructed conveniently as

$$\mathbf{3}_P := \Sigma P + \mathbf{1},$$

where ΣP is the *suspension* of P ¹. The two points of the suspension are called \mathbf{N} and \mathbf{S} , and the identity path (if it exists) between those points is called $\text{merid}(p) : \mathbf{N} = \mathbf{S}$, with $p : P$.

Recall the following about suspensions.

¹The suspension of a type is not generally a quotient, because it is not generally a set: we use the fact that P is a proposition here.

- By induction, we can define a map

$$\text{swap} : \Sigma P \rightarrow \Sigma P$$

that sends \mathbf{N} to \mathbf{S} and vice versa.

- By induction, we can define a map $\text{extract} : \mathbf{N} =_{\Sigma P} \mathbf{S} \rightarrow P$, and this can be generalized to a map

$$\text{extract}'_x : x =_{\Sigma P} \text{swap}(x) \rightarrow P.$$

Notice that if we have P , then the suspension is contractible, so $\mathbf{3}_P \simeq \mathbf{2}$, and also $(\mathbf{3}_P \simeq \mathbf{3}_P) \simeq \mathbf{2}$.

Define

$$g := f_{\mathbf{3}_P \simeq \mathbf{3}_P}(\text{ide}_{\mathbf{3}_P}) : \mathbf{3}_P \simeq \mathbf{3}_P,$$

where $\text{ide}_{\mathbf{3}_P}$ is the equivalence $\mathbf{3}_P \simeq \mathbf{3}_P$ given by the identity function on $\mathbf{3}_P$. We will see g both as an equivalence and as a function $\mathbf{3}_P \rightarrow \mathbf{3}_P$.

Now we do case analysis on $g(\text{inr}(\star))$. Notice that this case analysis is simply an instance of the induction principle for sum types. In particular, we do not require decidable equality of $\mathbf{3}_P$ (which would already give us $P + \neg P$, which is exactly what we are trying to prove). When analyzing the case $\text{inr}(t) : \mathbf{3}_P$, with $t : \mathbf{1}$, we are free to specialize to $t = \star$ since $\mathbf{1}$ is contractible.

$g(\text{inr}(\star)) = \text{inr}(\star)$: Assume that P holds. Then by transporting the witness of $f_{\mathbf{2}}(0_{\mathbf{2}}) \neq 0_{\mathbf{2}}$ along an equivalence that identifies $0_{\mathbf{2}}$ with $\text{ide}_{\mathbf{3}_P}$, we get that $g \neq \text{ide}_{\mathbf{3}_P}$. However, since $\mathbf{3}_P \simeq \mathbf{2}$ and g has a fixed point $\text{inr}(\star)$, we can deduce that $g = \text{ide}_{\mathbf{3}_P}$, which is a contradiction.

$g(\text{inr}(\star)) = \text{inl}(x)$: We do further case analysis on $g(\text{inl}(x))$.

$g(\text{inl}(x)) = \text{inr}(\star)$: We do further case analysis on $g(\text{inl}(\text{swap}(x)))$.

$g(\text{inl}(\text{swap}(x))) = \text{inr}(\star)$: Since we now have

$$g(\text{inl}(x)) = \text{inr}(\star) = g(\text{inl}(\text{swap}(x)))$$

and since g is an equivalence, we can use $\text{extract}'_x$ to get P .

$g(\text{inl}(\text{swap}(x))) = \text{inl}(y)$: Assume P , in which case $x = \text{swap}(x)$. Hence $\text{inr}(\star) = \text{inl}(y)$ which is a contradiction.

$g(\text{inl}(x)) = \text{inl}(y)$: Assume P , in which case $\text{inl}(x) = \text{inl}(y)$. But we now have

$$g(\text{inr}(\star)) = \text{inl}(x) = \text{inl}(y) = g(\text{inl}(x)).$$

So since g is an equivalence, this yields $\text{inr}(\star) = \text{inl}(x)$, which is a contradiction. □

Acknowledgements

Thanks to Martín Escardó, my supervisor, for his support. Thanks to Uday Reddy for giving the talk on parametricity that inspired me to think about this.

References

- [1] A. B. Booij. *Parametricity and excluded middle in Agda*. University of Birmingham, UK. URL: <http://www.cs.bham.ac.uk/~abb538/agda/nonparametric.html>.
- [2] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. Institute for Advanced Study: <http://homotopytypetheory.org/book>, 2013.